



Shrimp survival and growth can degrade quickly when primary stocks are reproduced as secondary stocks.

Summary:

Shrimp breeders often protect their genetic investment by selling broodstock that will accumulate ever-increasing levels of inbreeding in successive generations. The general procedure is widely known, but using the formula below, the quantitative estimate of inbreeding effects on the survival of later generations can be surprisingly high, especially in inferior environments.

Shrimp Copyright: INBREEDING STRATEGIES EFFECTIVE AGAINST ILLEGAL COPYING OF GENETICALLY IMPROVED SHRIMP

out permission for any number of generations are called secondary stocks.

Protecting Primary Stocks

By understanding a few basic steps, shrimp breeders can protect their investment in their primary stocks by incorporating inbreeding strategies.

Step 1

The shrimp breeder chooses four of the best mature shrimp from the broodstock to be parents of the production breeders in the following year. The selected shrimp should be best both in terms of their own phenotype and the known genetic quality of their near relatives, as well. Identifying the “best” animals constitutes 90% of the art of aquaculture genetics. As an illustration, four primary breeders, two males and two females, are colored blue and placed at the top of the pedigree diagram in Figure 1.

Step 2

One pair of blue primary breeders is used to produce thousands or tens of thousands of brothers, while the other pair produces a similar number of sisters. These two set of siblings are matured to become the production breeders.

Although each pair of primary breeders provides offspring of only one sex to the protection scheme illustrated in Figure 1, the offspring of the other sex need not be wasted. They can be used to produce another line of protected breeders.

Sometimes production breeders are sold to customers, and sometimes the PLs produced by mating them are sold. Whichever stage is sold, no inbreeding is added.

Step 3

The customer of the primary breeding company grows the PLs or mates breeders he purchased. Inbreeding does not occur in the shrimp grown to harvest. All is well.

Roger W. Doyle, Ph.D.
Genetic Computation Ltd.
4630-1 Lochside Drive
Victoria, British Columbia, Canada V8Y 2T1
rdoyle@genecomp.com

Dustin R. Moss, M.S.
Shaun M. Moss, Ph.D.
The Oceanic Institute
Waimanalo, Hawaii, USA

Shrimp breeders have invested a lot of money in the development of genetically improved broodstock because they want a bigger share of the world market for breeders and postlarvae (PLs). But how can they profit from their investment if their customers propagate the broodstock they buy, without permission? Or even worse, what can they do if their customers compete with them by selling postlarvae at a price that does not reflect the cost of genetic research and development?

Breeders often protect their investments in genetics through programs based on inbreeding. The general procedure is widely known, but using the formula below, the quantitative estimate of inbreeding effects on the survival of later generations can be surprising.

In the following, shrimp directly produced in genetic improvement programs are called primary stock. These include broodstock sold to multiplier hatcheries, as well as their offspring, which are sold to farmers as PLs for commercial growout. Broodstock and PLs that multiplier hatcheries produce by breeding primary stocks with-




Figure 1. The inbreeding level (F) of each animal is indicated in this pedigree of copy-protected shrimp. The four primary parents are colored blue. Males are denoted by squares, females by circles. Primary stocks are bracketed with "P." Those sold as breeders (row 2) or postlarvae (row 3) for growout are bracketed with "C." Primary stocks propagated without permission (secondary stocks) are bracketed with "S."

Inbred Secondary Stocks

Unauthorized reproduction of primary animals as secondary stocks leads to progressive inbreeding, as described below.

Step 1

The customer grows and then matures PLs purchased from the primary hatchery or produced by the purchased primary breeders, and mates them among themselves to generate a secondary stock (row 4 in Figure 1). Inbreeding, denoted F , will be either 0.125 or 0.25 in the first generation of a secondary stock.

The proportion of the two inbreeding levels in the secondary stock depends on the number of secondary spawners and their family relationships. The customer is now a secondary producer. He has no family relationship information to use for minimizing inbreeding in the shrimp and doesn't know the inbreeding level of any particular family of shrimp.

Step 2

Secondary spawning continues for more generations and inbreeding increases quickly, as shown in Figure 1. Again, the proportion of families with a given level of inbreeding varies with the numbers of spawners and relationship information, which is not available. An inbreeding level of $F = 0.5$ (equivalent to self-fertilization) is reached by some families in the third generation in a well-planned broodstock protection scheme. Inbreeding continues to accumulate in later generations.

Inbreeding Depression, Copy Protection

The F value of inbreeding is not in itself enough to help people make decisions. What everyone wants to know is the extent to which any given value of inbreeding depresses growth, reproductive performance, and survival in secondary stocks.




Figure 2. Relationship between inbreeding depression and environmental quality. Each data point summarizes an experiment at the Oceanic Institute which measured the survival of *L. vannamei* with various levels of inbreeding. The regression coefficient of survival on inbreeding within each experiment is shown on the Y axis as F -sensitivity. The projected survival at zero inbreeding is shown on the X axis. The curved red line is a logarithmic regression in which the sources of mortality are multiplicative rather than additive.

Workers at the Oceanic Institute in Hawaii, USA have found that inbreeding depression in Pacific white shrimp, *Litopenaeus vannamei*, depends on the quality of the environment. Environmental quality can be quantitatively characterized in this context as the survival of noninbred shrimp.

Data show that the worse the environment – i.e., the lower the survival of noninbred shrimp – the larger the depressive effect on inbred shrimp. The depression of growth rate and reproductive performance appears to be qualitatively similar to survival, but possibly somewhat less for growth and more for reproduction.

The magnitude of the effect is shown in Figure 2, where inbreeding depression is represented on the vertical axis by the variable F -sensitivity. F -sensitivity is the slope of the linear relationship between survival and inbreeding in a particular environment. The quality of the environment is measured on the horizontal axis by the survival rate of noninbred shrimp. When the environment is poor, and noninbred shrimp have low survival, the effect on the survival of inbred shrimp is strongly negative, i.e. they are very sensitive to inbreeding.

The following is a summary of F -sensitivity estimates in environments where survival of noninbred shrimp ranged from high to low because of experimental variations in salinity, challenges with viral pathogens, etc.

WHEN THE ENVIRONMENT IS POOR, THE EFFECT ON THE SURVIVAL OF INBRED SHRIMP IS STRONGLY NEGATIVE.

Predicting Broodstock Protection

If the survival of the primary stock where inbreeding is zero is 70%, what should the survival of a family with inbreeding $F = 0.25$ be in the same environment? The thin black lines in Figure 2 illustrate how to use this diagram to predict the effectiveness of copy protection.

Reading upward from 0.7 (70% survival) on the horizontal axis to the curved regression line, then sideways back to the vertical axis, we see that the F -sensitivity in that environment is about -0.65. Survival of secondary stocks with $F = 0.25$ can now be estimated from the formula:

$$\text{Secondary Survival} = \text{Primary Survival} + (F\text{-sensitivity} \times F).$$

Thus, in the above illustration, secondary survival is expected to equal $0.70 + (-0.65 \times 0.25) = .5375$ or about 54% – 16% lower than the noninbred shrimp. This may not be enough to worry about. But when the environment deteriorates and survival of the primary stocks falls, the results are not so good.

If primary stock survival drops to 40%, secondary stocks with 0.25 inbreeding would have $0.4 + (-1.3 \times 0.25) = 8\%$ survival, a serious problem. At primary stock survival of 50%, the survival of secondary stocks with $F = 0.375$ would be about 11%, while secondaries with 0.5 inbreeding would face total mortality.

The effect of broodstock protection is fairly clear in these examples. When the environment is good and the survival of noninbred shrimp is greater than 50%, inbreeding depression won't cause much of a problem until the third generation of illegal production. But in times of stress, when dissolved-oxygen concentration, water temperature, disease, or some combination of these factors

reduces primary survival, secondary stocks are at a high risk of being wiped out in one or two generations.

Controlling Multiplier Hatcheries

The broodstock protection scheme in Figure 1 is very simple, and additional protective steps can be taken. For instance, if a secondary breeder (multiplier hatchery) accumulates several batches from the primary stock and treats them as separate "families" to escape inbreeding, a canny primary breeder need merely create appropriate family relationships among the primary parents of successive batches.

These extra relationships will cause secondary inbreeding if several multiplier hatcheries band together to crossbreed primary stocks purchased in separate batches. Of course, these protective steps add a bit to the operating cost of the genetic improvement program, but not very much in a program that is set up properly.

If secondary breeders hybridize protected shrimp with stocks from an entirely different population to avoid inbreeding, the initial secondary generation will be non-inbred, but inbreeding will accumulate quickly after that because of the close family relationship among one-half of the parents in the hybrid cross.

Genetic Traps Effective

There is not a lot secondary breeders can do to escape the genetic traps set by vigilant primary breeders. The best strategy for multiplier hatcheries is to purchase the latest generation of improved broodstock every year. For hatcheries unwilling to do that, the authors' best advice is to spawn as many breeders as possible, keep track of what generation of illegal PLs is being produced, and hope for really nice weather in generation 4.